

**ASTHMA REFERRAL FORM**

Please complete the following and fax with clinical documentation to: 720.279.7461

Referral Process			
<b>1. PATIENT INFORMATION</b>		<b>2. PHYSICIAN INFORMATION</b>	
Name:		Physician's name:	
Address:		License #:                      NPI #:	
City:	State:                      Zip:	DEA #:	
Home Phone:	Other Phone:	Address:	
Email:		City:                      State:                      Zip:	
DOB:	Social Security #:	Office Contact:	
Gender: <input type="checkbox"/> M <input type="checkbox"/> F	Height:                      Weight:	Phone:                      Fax:	
Allergies:			
<b>3. DIAGNOSIS</b> Year of diagnosis: _____ <input type="checkbox"/> Moderate Persistent Asthma (J45.4) <input type="checkbox"/> Severe Persistent Asthma (J45.5) <input type="checkbox"/> Chronic Idiopathic Urticaria (L50.1) <input type="checkbox"/> Chronic Rhinosinusitis (J32.9) with Nasal Polyposis (J33.9) <input type="checkbox"/> Severe persistent asthma with (acute) exacerbation (J45.51) <input type="checkbox"/> Other(s): _____			
<b>4. INSURANCE INFORMATION: Please submit copies of the front and back of primary and secondary insurance cards with this referral.</b>			
<b>5. ADDITIONAL INFORMATION REQUESTED</b> Prescription type: <input type="checkbox"/> Naive/new start <input type="checkbox"/> Restart <input type="checkbox"/> Continued therapy  Lab results: <input type="checkbox"/> History of positive skin OR RAST test to a perennial aeroallergen    IgE Level: _____ IU/mL; date _____  Current/previous history of parasitic (helminth) infections? _____  Concomitant Therapies: <i>check all that apply:</i> <input type="checkbox"/> Short acting beta agonist <input type="checkbox"/> Long acting beta agonist <input type="checkbox"/> Antihistamines <input type="checkbox"/> Decongestants <input type="checkbox"/> Immunotherapy <input type="checkbox"/> Inhaled Corticosteroid <input type="checkbox"/> Leukotriene modifiers <input type="checkbox"/> Oral steroid <input type="checkbox"/> Nasal steroid <input type="checkbox"/> Other: _____			
<b>6. PRESCRIPTION INFORMATION</b>  <input type="checkbox"/> <b>Fasenra™ (benralizumab) 30 mg/1 ml pre-filled syringe</b>  Inject 30 mg (1ml) subcutaneously every 4 weeks for the first 3 doses, followed by 30 mg once every 8 weeks thereafter		<b>Quantity</b> _____ syringes <input type="checkbox"/> 30-day supply <input type="checkbox"/> 90-day supply <input type="checkbox"/> _____-day supply	<b>Refills</b> <input type="checkbox"/> 1 year <input type="checkbox"/> _____
<input type="checkbox"/> <b>CINQAIR® (reslizumab) – 100 mg/10mL vial</b>  Inject 3 mg/kg once every 4 weeks by IV infusion over 20 to 50 minutes		<b>Quantity</b> _____ vials <input type="checkbox"/> 30-day supply <input type="checkbox"/> 90-day supply <input type="checkbox"/> _____-day supply	<b>Refills</b> <input type="checkbox"/> 1 year <input type="checkbox"/> _____

<p><b>Asthma Adult Dosing</b></p> <p><input type="checkbox"/> <b>Dupixent® (dupilumab) 200 mg/1.14 ml prefilled syringe</b> Initial dose of 400 mg (two 200 mg injections) subcutaneously followed by 200 mg every other week</p> <p><input type="checkbox"/> <b>Dupixent® (dupilumab) 300 mg/2 ml prefilled syringe</b> Initial dose of 600 mg (two 300 mg injections) subcutaneously followed by 300 mg every other week</p> <p><b>CRSwNP</b></p> <p><input type="checkbox"/> <b>Dupixent® (dupilumab) 300 mg/2 ml prefilled syringe</b> Recommended dose is 300 mg subcutaneously every other week</p> <p><b>Asthma Pediatric Dosing</b> <i>For 15 to less than 30 kg:</i></p> <p><input type="checkbox"/> 100 mg subcutaneously every 2 weeks</p> <p><b>OR</b></p> <p><input type="checkbox"/> 300 mg subcutaneously every 4 weeks</p> <p><i>For ≥ 30 kg:</i> <input type="checkbox"/> 200 mg subcutaneously every 2 weeks</p>	<p><b>Quantity</b></p> <p>_____ syringes</p> <p><input type="checkbox"/> 30-day supply</p> <p><input type="checkbox"/> 90-day supply</p> <p><input type="checkbox"/> _____-day supply</p>	<p><b>Refills</b></p> <p><input type="checkbox"/> 1 year</p> <p><input type="checkbox"/> _____</p>
<p><input type="checkbox"/> <b>Nucala® (mepolizumab) – 100 mg vial</b> Inject 100 mg subcutaneously once every 4 weeks into the upper arm, thigh or abdomen</p> <p>Include sterile water and supplies sufficient for medication administration</p> <ul style="list-style-type: none"> <li>• One 10 mL vial sterile water for injection for every vial of Nucala® dispensed</li> <li>• Alcohol swabs</li> <li>• 3 mL Luer Lock injection syringe</li> <li>• NDL 21G needle for reconstitution</li> <li>• 1 mL polypropylene syringe with 21G to 27G x ½" needle for subcutaneous injection</li> </ul> <p><input type="checkbox"/> No supplies</p>	<p><b>Quantity</b></p> <p><input type="checkbox"/> 30-day supply</p> <p><input type="checkbox"/> 90-day supply</p> <p><input type="checkbox"/> _____-day supply</p>	<p><b>Refills</b></p> <p><input type="checkbox"/> 1 year</p> <p><input type="checkbox"/> _____</p>
<p><input type="checkbox"/> <b>Xolair® (omalizumab) prefilled syringe (<i>dispense minimal number of 75mg and/or 150mg to complete prescribed dose</i>)</b></p> <p><b>OR</b></p> <p><input type="checkbox"/> <b>Xolair® (omalizumab) 150mg single dose vial</b></p> <p><b>Every 4 weeks dosing:</b></p> <p><input type="checkbox"/> Administer 75mg per dose subcutaneously every 4 weeks</p> <p><input type="checkbox"/> Administer 150mg per dose subcutaneously every 4 weeks</p> <p><input type="checkbox"/> Administer 225mg per dose subcutaneously every 4 weeks</p> <p><input type="checkbox"/> Administer 300mg per dose subcutaneously every 4 weeks</p> <p><input type="checkbox"/> Administer 450mg per dose subcutaneously every 4 weeks</p> <p><input type="checkbox"/> Administer 600mg per dose subcutaneously every 4 weeks</p> <p><input type="checkbox"/> Administer other: _____ mg per dose subcutaneously every 4 weeks</p> <p><b>Every 2 weeks dosing:</b></p> <p><input type="checkbox"/> Administer 225mg per dose subcutaneously every 2 weeks</p> <p><input type="checkbox"/> Administer 300mg per dose subcutaneously every 2 weeks</p> <p><input type="checkbox"/> Administer 375mg per dose subcutaneously every 2 weeks</p> <p><input type="checkbox"/> Administer 450mg per dose subcutaneously every 2 weeks</p> <p><input type="checkbox"/> Administer 525mg per dose subcutaneously every 2 weeks</p> <p><input type="checkbox"/> Administer 600mg per dose subcutaneously every 2 weeks</p> <p><input type="checkbox"/> Administer other: _____ mg per dose subcutaneously every 2 weeks</p>	<p>Xolair® (omalizumab)</p> <p><input type="checkbox"/> Asthma (dose is dependent on weight and IgE levels, see package insert)</p> <p><input type="checkbox"/> CIU (fixed dose, not dependent on weight or IgE)</p> <p><input type="checkbox"/> Nasal Polyposis</p>	<p><b>Quantity</b></p> <p><input type="checkbox"/> 30-day supply</p> <p><input type="checkbox"/> 90-day supply</p> <p><input type="checkbox"/> _____-day supply</p> <p><b>Refills</b></p> <p><input type="checkbox"/> 1 year</p> <p><input type="checkbox"/> _____</p>

**FLUSH ORDERS**

**PIV/midline/PICC:** Flush before, after each infusion, and as needed with 3-20 ml NS, followed by Heparin 2-5 ml if indicated

Heparin 10 unit/ml       Heparin 100 unit/ml

**Port:** Flush before, after each infusion, and as needed with 5-20 ml NS, followed by Heparin 100 unit/ml 5 ml.

**ANAPHYLAXIS ORDERS**

**Adults:** For mild reaction-administer diphenhydramine 50mg IVP/PO x 1.

For severe reactions-administer epinephrine 0.3mg SubQ/IM x 1. May repeat x1 if needed. Contact 911 and monitor vital signs.

Other: \_\_\_\_\_

**Pediatrics:** administer by age: *For mild reaction (rash/hives) give diphenhydramine*

**Age 1-5:** 12.5mg IV/PO x1      **Age 6-11:** 25mg IV/PO x1      **Age 12+:** 50mg IV/PO x1

*For severe reaction (airway closure) administer epinephrine. 0.01mg/kg/dose (max 0.3mg) SubQ/IM x 1.*

*If needed, repeat a second dose after 5-10 minutes for a maximum of two doses. Contact 911 and monitor vital signs.*

7.  Dispense as written     Substitution Permitted

PHYSICIAN'S SIGNATURE (required): \_\_\_\_\_ Date: \_\_\_\_\_

[www.veroshealth.com](http://www.veroshealth.com)

© Veros Health. All rights reserved. 01.22

Prescription is valid for one year unless otherwise indicated. The prescriber is to comply with his/her state specific prescription requirements such as e-prescribing, state specific prescription form, fax language, etc. Non-compliance with state specific requirements could result in outreach to the prescriber.