

GI Referral Form

Please complete the following and fax with clinical documentation to:
720.279.7461

Referral Process	
1. PATIENT INFORMATION	2. PHYSICIAN INFORMATION
Name: _____	Physician's name: _____
Address: _____	License #: _____ NPI #: _____
City: _____ State: _____ Zip: _____	DEA #: _____
Home Phone: _____ Other Phone: _____	Address: _____
Email: _____	City: _____ State: _____ Zip: _____
DOB: _____ Social Security #: _____	Office Contact: _____
Gender: <input type="checkbox"/> M <input type="checkbox"/> F Height: _____ Weight: _____	Phone: _____ Fax: _____
3. DIAGNOSIS <input type="checkbox"/> Crohn's (K50.00) <input type="checkbox"/> Ulcerative Colitis (K51.90) <input type="checkbox"/> Other _____ TB Test? <input type="checkbox"/> No <input type="checkbox"/> Yes If yes, result: <input type="checkbox"/> Positive <input type="checkbox"/> Negative	
4. INSURANCE INFORMATION Please submit copies of the front and back of primary and secondary insurance cards with this referral.	
5. PRESCRIPTION INFORMATION (Medication, Dosage, Duration, Pre-Medication)	
Cimzia® <input type="checkbox"/> Loading dose: Inject 400 mg SC initially and at weeks 2 and 4. # of Refills _____ <input type="checkbox"/> Inject 400 mg SC every _____ weeks. # of Refills _____ Simponi® <input type="checkbox"/> Initial dose: Inject 200 mg SC at week 0, followed by 100 mg at week 2 <input type="checkbox"/> Maintenance doses: Inject 100 mg SC every 4 weeks. # of Refills _____ Stelara® 130 mg/26 mls (5mg/ml) single dose vials for intravenous use Initial IV infusion: <input type="checkbox"/> ≤55kg 260 mg IV over 1 hour x 1 dose <input type="checkbox"/> 55kg to 85 kg 390 mg IV over 1 hour x 1 dose <input type="checkbox"/> >85kg 520 mg IV over 1 hour x 1 dose Stelara® for subcutaneous use <input type="checkbox"/> 45 mg/0.5mL solution in a single-dose vial <input type="checkbox"/> 90 mg/1 ml Prefilled Syringe <input type="checkbox"/> Maintenance: Inject 90mg (1 ml) SC 8 weeks after initial IV infusion then every 8 weeks thereafter. Refills x _____ Months Entyvio® <input type="checkbox"/> 300 mg infused intravenously over approximately .30 minutes at 0, 2 and 6 weeks, then every 8 weeks thereafter Humira® <input type="checkbox"/> Initially - Inject 160mg subcutaneously day one; 80mg day 15; 40mg day 29 <input type="checkbox"/> Maintenance Dose - Inject 40mg subcutaneously every other week Injection training required? <input type="checkbox"/> Yes <input type="checkbox"/> No Remicade® <input type="checkbox"/> Loading dose: 5mg/kg @ 0, 2 and 6 weeks <input type="checkbox"/> Infuse in NS 250ml over 2 hours as directed <input type="checkbox"/> Maintenance: 5mg/kg q 8 week Maintenance Dose. Refills x _____ Months. <input type="checkbox"/> Other _____	
6. ANAPHYLAXIS ORDERS	
Adults: For mild reaction-administer diphenhydramine 50mg IVP/PO x 1. For severe reactions-administer epinephrine 0.3mg SubQ/IM x 1. May repeat x1 if needed. Contact 911 and monitor vital signs. Pediatrics: administer by age: For mild reaction (rash/hives) give diphenhydramine Age 1-5: 12.5mg IV/PO x1 Age 6-11: 25mg IV/PO x1 Age 12+: 50mg IV/PO x1 For severe reaction (airway closure) administer epinephrine. 0.01mg/kg/dose (max 0.3mg) SubQ/IM x 1. If needed, repeat a second dose after 5-10 minutes for a maximum of two doses. Contact 911 and monitor vital signs. <input type="checkbox"/> Other: _____	
7. <input type="checkbox"/> Dispense as written <input type="checkbox"/> Substitution Permitted	
PHYSICIAN'S SIGNATURE (required): _____ Date: _____	