

Soliris® Referral Form

Please complete the following and fax with clinical documentation to: 720.279.7461

Referral Process

Referral Process	
1. PATIENT INFORMATION	2. PHYSICIAN INFORMATION
Name: _____	Physician's name: _____
Address: _____	License #: _____ NPI #: _____
City: _____ State: _____ Zip: _____	DEA #: _____
Home Phone: _____ Other Phone: _____	Address: _____
Email: _____	City: _____ State: _____ Zip: _____
DOB: _____ Social Security #: _____	Office Contact: _____
Gender: <input type="checkbox"/> M <input type="checkbox"/> F Height: _____ Weight: _____	Phone: _____ Fax: _____
3. DIAGNOSIS Year of diagnosis: _____	
<input type="checkbox"/> Myasthenia Gravis w/o acute exac. (G70.00) <input type="checkbox"/> Myasthenia Gravis with acute exac. (G70.01) ACHR-AB results (required) _____ MGFA Classification _____ MG-ADL Scores _____	<input type="checkbox"/> Neuromyelitis Optica (G36.0) AQP4 Autoantibodies test (required) _____ MRI _____ CSF test _____
4. INSURANCE INFORMATION Please submit copies of the front and back of primary and secondary insurance cards with this referral.	
5. ADDITIONAL INFORMATION REQUESTED Please provide relevant history, including office and hospitalization notes	
Allergies: _____	
Patient has received meningococcal vaccine <input type="checkbox"/> Yes <input type="checkbox"/> No Date Received _____	
Previous tried and failed meds (immune suppressants, IVIG, plasma exchange): _____	
PRESCRIPTION INFORMATION Anticipated Start Date: _____	
Soliris® (eculizumab)	
<ul style="list-style-type: none"> 900 mg weekly for the first 4 weeks, followed by 1200 mg for the fifth dose 1 week later, then 1200 mg every 2 weeks thereafter 	
6. FLUSH ORDERS	
PIV/midline/PICC: Flush before, after each infusion, and as needed with 3-20 ml NS, followed by Heparin 2-5 ml if indicated <input type="checkbox"/> Heparin 10 unit/ml <input type="checkbox"/> Heparin 100 unit/ml	
Port: Flush before, after each infusion, and as needed with 5-20 ml NS, followed by Heparin 100 unit/ml 5 ml.	
ANAPHYLAXIS ORDERS	
Adults: For mild reaction-administer diphenhydramine 50mg IVP/PO x 1. For severe reactions-administer epinephrine 0.3mg SubQ/IM x 1. May repeat x1 if needed. Contact 911 and monitor vital signs.	
Skilled Nursing Services Needed? <input type="checkbox"/> Yes <input type="checkbox"/> No Additional Instructions: _____	
Dispense quantity: _____ Refills: _____	
6. <input type="checkbox"/> Dispense as written <input type="checkbox"/> Substitution Permitted	
PHYSICIAN'S SIGNATURE (required): _____ Date: _____	